

MEAN CURVATURE MOTION OF TRIPLE JUNCTIONS OF GRAPHS IN TWO DIMENSIONS

by A. Freire

University of Tennessee, Knoxville

Abstract

We consider a system of three surfaces, graphs over a bounded domain in $\mathbb{R}^2$, intersecting along a time-dependent curve and moving by mean curvature while preserving the pairwise angles at the curve of intersection (equal to $2\pi/3$.) For the corresponding two-dimensional parabolic free boundary problem we prove short-time existence of classical solutions (in parabolic Hölder spaces), for sufficiently regular initial data satisfying a compatibility condition.

0. Introduction: Triple junctions of graphs.

The goal of this paper is to prove short-time existence for the following geometric evolution problem: a configuration of three surfaces in $\mathbb{R}^3$, bordered by a common time-dependent curve of intersection $\Lambda(t)$, moves with normal velocity given by mean curvature (at each interior point) in such a way that the pairwise angles defined by their unit normals along $\Lambda(t)$ are constant throughout the evolution, equal to $2\pi/3$ radians.

The corresponding evolution for systems of curves was considered in [BronsardReitich] (where short-time existence was proved) and in [Mantegazza et al.], which includes a continuation criterion and the blow-up analysis at the first singular time. Very recently, the preprint [LensSeminar] includes global existence results for symmetric systems of curves of ‘lens type’. In addition to being the natural time-dependent version of the classical problem of minimal surfaces meeting along a ‘liquid edge’ ([Dierkes et al.] p. 299 ff.), the problem derives its interest from being the sharp-interface limit of a well-known class of parabolic evolutionary models in materials science, defined by a vector-valued order parameter and a multi-well potential. (See [BronsardReitich] for a discussion in the case of curves.)

In the case of surfaces, the parametric approach adopted to prove local existence for curves in [BronsardReitich] and [Mantegazza et al.] does not work, that is, does not lead directly to a well-posed parabolic system. The essential difficulty can be traced back to the fact that the junction is now a one-dimensional object, and one does not expect that independently evolving parametrizations of each surface in a triple junction configuration will continue to match pointwise along the junction, even if they do so at

$t = 0$. On the other hand, it is difficult to encode the matching condition analytically so as to allow for such ‘sliding’ along the junction. Thus one is naturally led to consider the special case of *graphs*, for the simple reason that the graph parametrization is canonical, so that the surfaces’ meeting along $\Lambda(t)$ corresponds to three functions coinciding pointwise along a moving interface Γ_t in $\mathbb{R}^2$.

In this way the problem for graphs becomes a more-or-less classical free boundary problem for a quasilinear parabolic system in two dimensions. By analogy with the usual Stefan problem terminology, this would be a ‘three-phase problem’, with two of the ‘phases’ coexisting in the same domain. The corresponding ‘one-phase’ problem- the mean curvature motion of a single graph over a time-dependent domain $D(t)$ in $\mathbb{R}^n$, intersecting $\mathbb{R}^n$ at a constant angle- has been dealt with in a recent preprint of the author ([Freire]), including local existence and some results on long-time behavior. In [Freire], we introduced additional ‘orthogonality’ conditions at the free boundary to obtain a well-posed system. If one tries to extend this method to the more general case considered here, compositions of the unknown functions with the (also unknown) diffeomorphisms pulling back the evolution to a fixed domain come into play, and since compositions behave poorly in Hölder spaces it becomes difficult to implement a fixed-point scheme using this method. Instead, we use here an ingenious transformation method which we learned from the paper [BaconneauLunardi]. (These authors considered a one-phase multidimensional free-boundary problem for a semilinear parabolic system).

Now for a precise description of the problem and results. We focus in this paper on configurations parametrized by two disks and one annulus, but it is clear the method would work for more general configurations (for example, two annuli over one sub-domain, one disk over the other; or three annuli.) In addition, the system lives in a fixed bounded domain $\Omega \subset \mathbb{R}^2$, and we must specify how one of the surfaces intersects $\partial\Omega$; we choose orthogonal intersection, which leads to a standard Neumann condition at the fixed boundary. More generally, one could prescribe orthogonal intersection with a non-vertical, fixed ‘support surface’. This would introduce additional moving boundaries into the problem but analytically would present no essential difficulty, so we avoided this in the interest of simplicity.

A *configuration* at time t consists of three surfaces:

$$\Sigma_t^I = \text{graph}(w_t^I), \quad w_t^I = w^I(\cdot, t) : D_t^I \rightarrow \mathbb{R}, \quad I = 1, 2, 3,$$

where $D_t^1 = D_t^2$ and $D_t^3 = \Omega - \bar{D}_t^1$ are bounded, connected open subsets of the bounded domain $\Omega \subset \mathbb{R}^2$. The Σ_t^I intersect along an embedded closed

curve $\Lambda_t \subset \mathbb{R}^3$, a graph over the simple closed planar curve $\Gamma_t = \bar{D}_t^1 \cap \bar{D}_t^3$. The upper unit normals are:

$$N^I = \frac{1}{v^I}[-Dw^I, 1], \quad v^I = \sqrt{1 + |Dw^I|^2}.$$

Along Γ_t , we impose the following conditions:

- (i) (matching) $w^1(y, t) = w^2(y, t) = w^3(y, t), \quad y \in \Gamma_t \quad (BC_w^0).$
- (ii) (angle) $N^1 + N^2 = N^3.$

That is, the Σ^I intersect making pairwise $2\pi/3$ angles. In addition, Σ^3 intersects the vertical cylinder over $\partial\Omega$ orthogonally: if ω denotes the inner unit normal to $\partial\Omega$, we require:

$$d_\omega w^3 = 0 \text{ on } \partial\Omega. \quad (BC_{w^3})$$

The angle condition is equivalent to two scalar conditions on Γ_t :

$$\begin{cases} \frac{1}{v^1} + \frac{1}{v^2} = \frac{1}{v^3}, & (BC_w^1) \\ \frac{d_n w^1}{v^1} + \frac{d_n w^2}{v^2} = \frac{d_n w^3}{v^3}, & (BC_w^2) \end{cases}$$

where n is the inner unit normal of Γ_t .

To specify a configuration, we need three time-dependent functions $w^I(t) : D_t^I \rightarrow \mathbb{R}$, defining graph parametrizations of the Σ_t^I :

$$G^I : E_T^I \rightarrow \mathbb{R}^3, \quad E_T^I = \{(x, t) \in \mathbb{R}^2 \times [0, T]; x \in D_t^I\}$$

$$G^I(x, t) = [x, w^I(x, t)] \in \mathbb{R}^2 \times \mathbb{R}.$$

(The labeling of the surfaces is chosen so that near $\Lambda(t)$, Σ^2 lies below Σ^1 .)

We wish to evolve a given initial configuration through configurations, so that on each surface the normal velocity is given by the mean curvature at each point:

$$\langle \partial_t G^I, N^I \rangle = H^I.$$

Recall $H^I = \text{tr}_{g^I} h^I = g^{Iij} h_{ij}^I$, where h^I is the second fundamental form of Σ^I , pulled back to $\mathbb{R}^2$ via G :

$$h^I(e, e) = -\langle d_e N^I, d_e G^I \rangle = \langle d^2 G^I(e, e), N^I \rangle = \frac{1}{v^I} d^2 w^I(e, e), \quad e \in \mathbb{R}^2,$$

and g^I is the induced metric:

$$g_{ij}^I = \langle \partial_i G^I, \partial_j G^I \rangle = \delta_{ij} + \partial_i w^I \partial_j w^I.$$

Mean curvature motion of graphs is given by the equation:

$$(MCM)_w \quad L_{g^I}[w^I] = 0, \quad L_{g^I}[f] := \partial_t f - \text{tr}_{g^I} D^2 f.$$

Our local existence result follows:

Theorem. *Let $\Omega \subset \mathbb{R}^2$ be a bounded open set with $C^{3+\alpha}$ boundary, $D_0^1 = D_0^2$ and $D_0^3 = \Omega - \bar{D}_0^1$ open subsets of Ω with common boundary component Γ_0 , a simple closed $C^{3+\alpha}$ curve in Ω . Given $w_0^I \in C^{3+\alpha}(D_0^I)$, assume:*

- (1) *The $\Sigma_0^I = \text{graph}(w_0^I)$ define a triple junction configuration;*
- (2) *The mean curvatures H_0^I of Σ_0^I satisfy at points of Γ_0 the following compatibility condition:*

$$H_0^1 + H_0^2 = H_0^3, \quad y \in \Gamma_0.$$

Then there exists $T > 0$ depending only on the initial data, domains $E_T^I \subset \Omega \times [0, T]$ and functions $w^I \in C^{2+\alpha, 1+\alpha/2}(E_T^I)$ such that:

- (3) *w^I is a classical solution of $L_{g^I}[w^I] = 0$ in E^I ;*
- (4) *The conjugation conditions (BC_w) hold on $\Gamma_t = \partial D_t^1$, and the boundary condition (BC_{w^3}) holds on $\partial\Omega$.*

Notation.

$$E_T^I = \bigcup_{t \in [0, T]} D_t^I \times \{t\} \subset \Omega \times [0, T];$$

it is always assumed that $T = \sup\{t \in [0, T); D_t^I \neq \emptyset \forall I\}$. We use standard notation for parabolic Hölder spaces; for example, the norm in $C^{2+\alpha, 1+\alpha/2}(E)$ (where $E \subset \Omega \times [0, T]$) is given by:

$$\|f\|_{2+\alpha} := \|f\|_{C^{2,1}(E)} + [D^2 f]_x^{(\alpha)} + [D^2 f]_t^{(\alpha/2)} + [\partial_t f]_x^{(\alpha)} + [\partial_t f]_t^{(\alpha/2)} + [Df]_t^{((1+\alpha)/2)},$$

with the standard notations for Hölder-type difference quotients in x or in t . α is an arbitrary constant in the open interval $(0, 1)$, fixed throughout the paper.

The subscript t always denotes a map (or function, or set) at time t , never a partial derivative (denoted ∂_t). The unit vectors ν , τ_0 are the inner unit normal (resp. counterclockwise unit tangent) vectors to Γ_0 , while n and τ denote the corresponding unit vectors for Γ_t .

Remarks.

(1) Note the drop in regularity: we require $C^{3+\alpha}$ initial data for $C^{2+\alpha}$ (in space) solutions. This occurs also in the main result in [BaconneauLunardi], and for the same reason. It was also observed in [Freire], where local existence for the ‘one-phase’ problem is obtained by a different argument.

(2) The result, at this point, lacks both a uniqueness statement and a continuation criterion. It is conceivable that assuming higher regularity on the initial data ($C^{4+\alpha}$), a uniqueness statement could be obtained following the argument in [BaconneauLunardi]. Both issues will be considered in further work.

(3) The restriction to two dimensions makes the argument technically easier, but is probably inessential. On the other hand, generalizing to configurations beyond graphs may require a completely different line of argument.

Outline. Section 1: We derive the compatibility condition and the equation of motion for the junction $\Lambda(t)$ (and for the ‘interface’ Γ_t). Section 2: In a standard way, we use a time-dependent diffeomorphism to pass to a system (with *four* unknown functions) over the fixed domains D_0^I :

$$w^I(y, t), \quad y \in D_t^I \longrightarrow u^I(x, t), \tilde{\rho}(x, t), \quad x \in D_0^I,$$

where $\tilde{\rho}$ is an extension to Ω of the function ρ_t parametrizing Γ_t as a normal graph over Γ_0 . Section 3: Following the argument in [BaconneauLunardi], we introduce a transformation of the dependent variables:

$$u^I(x, t), \tilde{\rho}(x, t) \longrightarrow U^I(x, t), \quad x \in D_0^I,$$

which leads to a system on $\Omega \times [0, T]$ with three unknown functions and three conjugation conditions on Γ_0 . Section 4: We describe the fixed-point scheme, the associated linear problem and the required Hölder space estimates. This concludes the proof, except for verifying that the linear system satisfies the ‘complementarity conditions’, which is done in Section 5.

Acknowledgments. This research was started during a semester visit to the Max-Planck Institute for Gravitational Physics (Potsdam-Golm); I am grateful to the Geometric Analysis group for the invitation and to the Max-Planck Society for support. An early version of this result was announced at a seminar at the FU Berlin (June 2007), and the current version at the conference Nonlinear PDEs at IMPA (August 2008). Finally, it is a pleasure to thank N.Alikakos (University of Athens) for originally suggesting the problem and for his interest in the results.

1. Compatibility conditions.

We wish to find solutions that converge to the initial data in a strong sense, as $t \rightarrow 0_+$. It is well-known from parabolic theory that, for such solutions to exist, the initial data have to satisfy compatibility conditions; in the present case, of ‘order 0’ and ‘order 1’. Before going any further, one should check that the geometric problem is compatible with a graph formulation, in the sense that one can find reasonably large families of initial configurations satisfying these compatibility requirements. In this section we state the conditions, analyze their geometry and examine their internal consistency.

‘Order 0’ compatibility is just the requirement that the initial data satisfy the conjugation and boundary conditions (i.e., that the w_0^I define a configuration).

On Γ_0 this corresponds to two zero-order conditions on the w_0^I (pointwise matching) and two first-order conditions (angle). Assuming the $w_0^I|_{\Gamma_0}$ are given (and hence the curve Λ_0), the conditions on the first-order data (the tangent planes to $\Sigma^I(0)$ along $\Lambda(0)$) are clearly compatible. (Construction: at each point, on the plane orthogonal to the tangent line of Λ_0 - which is determined by zero-order data- we can arbitrarily pick one unit vector, say $N^1(0)$, and then the other two are uniquely determined; so we have one real degree of freedom.)

‘Order one’ compatibility conditions along Γ_0 are obtained by differentiating the incidence conditions in t and using the equation, bearing in mind that Γ is time-dependent.

Parametrizing Γ_t by $a(s, t) \in \mathbb{R}^2$ ($s \in \mathbb{R}/2\pi\mathbb{Z}$ is a fixed periodic variable, not arc length), we compute (using the equation of motion) the velocity vector at points of Λ_t , $V^I = \frac{\partial}{\partial t}(G^I(a(s, t), t))$:

$$V^I = [\dot{a}, Dw^I \cdot \dot{a} + \partial_t w^I] = [\dot{a}, Dw^I \cdot \dot{a} + v^I H^I],$$

where $\dot{a} = \partial_t a$.

The compatibility condition is then the statement that, at $t = 0$ (and, *a fortiori*, for all t):

$$V^1 = V^2 = V^3, \quad y \in \Gamma_t.$$

In principle, this corresponds to up to six independent scalar conditions. To analyze the situation, observe that at points of Λ_t we have three natural orthonormal frames $\{E, N^I, T^I\}$, where E is the unit tangent vector to Λ_t (orienting Γ_t counterclockwise) and we define $T^I = E \wedge N^I$ (vector product),

a tangent vector to Σ^I ; in particular, $T^1 + T^2 = T^3$. Noting that $\langle V^I, N^I \rangle = H^I$, we may write:

$$V^I = \lambda^I E + \mu^I T^I + H^I N^I.$$

Six independent scalar conditions equivalent to equality of the V^I are then obtained as follows:

- (1.) Add $\langle V^1, N^1 \rangle = \langle V^3, N^1 \rangle$ and $\langle V^2, N^2 \rangle = \langle V^3, N^2 \rangle$.
- (2.) Add $\langle V^1, T^1 \rangle = \langle V^3, T^1 \rangle$ and $\langle V^2, T^2 \rangle = \langle V^3, T^2 \rangle$.
- (3.) Take the difference $\langle V^1, N^1 \rangle = \langle V^3, N^1 \rangle$ minus $\langle V^2, N^2 \rangle = \langle V^3, N^2 \rangle$.
- (4.) Take the difference $\langle V^1, T^1 \rangle = \langle V^3, T^1 \rangle$ minus $\langle V^2, T^2 \rangle = \langle V^3, T^2 \rangle$.
- (5./6.) $\langle V^1, E \rangle = \langle V^2, E \rangle = \langle V^3, E \rangle$.

As we proceed to write these relations in a more explicit form, we will preserve their labels (1)...(6). We remark that conditions (1)...(4) already occur for curves, while (5) and (6) are new for surfaces.

In terms of the coefficients μ^I and λ^I , we easily obtain (using $N^1 + N^2 = N^3$ and $T^1 + T^2 = T^3$):

- (1) $H^1 + H^2 = H^3$.
- (2) $\mu^1 + \mu^2 = \mu^3$.
- (3) $H^1 - H^2 = \sqrt{3}\mu^3$.
- (Here we used $\langle T^3, N^1 - N^2 \rangle = \sqrt{3}$ and $\langle N^3, N^1 - N^2 \rangle = 0$.)
- (4) $\mu^1 - \mu^2 = -\sqrt{3}H^3$.
- (5/6) $\lambda^1 = \lambda^3$ and $\lambda^2 = \lambda^3$.

To make these relations explicit, one has to compute the T^I , λ^I and μ^I , which is elementary. Denote by τ the unit tangent vector to Γ_t , and set $v_E = \sqrt{1 + (d_\tau w^I)^2}$ (independent of I on Γ_t) and $(Dw^I)^\perp = (-\partial_2 w^I, \partial_1 w^I)$. The results are:

$$T^I = -\frac{1}{v^I v_E} [n - (d_\tau w^I)(Dw^I)^\perp, d_n w^I],$$

$$\mu^I = -\frac{1}{v_E} ((\dot{a} \cdot n)v^I + (d_n w^I)H^I),$$

$$\lambda^I = \frac{1}{v_E} (\dot{a} \cdot \tau + (d_\tau w^I)(\dot{a} \cdot Dw^I + v^I H^I)).$$

With this we can rewrite (1)-(4) as:

- (1) $H^1 + H^2 = H^3$.
- (2) $(\dot{a} \cdot n)(v^1 + v^2) + (d_n w^1)H^1 + (d_n w^2)H^2 = (\dot{a} \cdot n)v^3 + (d_n w^3)H^3$.
- (3) $H^1 - H^2 = (\sqrt{3}/v_E)((\dot{a} \cdot n)v^3 + (d_n w^3)H^3)$.
- (4) $(\dot{a} \cdot n)(v^1 - v^2) + (d_n w^1)H^1 - (d_n w^2)H^2 = -\sqrt{3}v_E H^3$.

When rewriting conditions (5/6), we observe that terms involving the tangential component $\dot{a} \cdot \tau$ of the moving boundary velocity cancel, and there is a common factor $d_\tau w \equiv d_\tau w^I$ (independent of I). We have:

$$\begin{aligned} (5/6) \quad (d_\tau w)((\dot{a} \cdot n)d_n w^1 + v^1 H^1) &= (d_\tau w)((\dot{a} \cdot n)d_n w^2 + v^2 H^2) \\ &= (d_\tau w)((\dot{a} \cdot n)d_n w^3 + v^3 H^3). \end{aligned}$$

Thus (5) and (6) occur only at points where $d_\tau w \neq 0$ (that is, where Λ_t is not ‘horizontal’). Regarding the zero and first order data of the configuration as given, (1)-(6) give six homogeneous linear relations for the four ‘second-order quantities’ $\dot{a} \cdot n$ and H^I , so it is conceivable that only the zero solution might exist. That this is not so is part of the conclusion of the following proposition.

Proposition 1.1. *The homogeneous system of linear relations (1)-(6) has rank two, and is generated by (1) and (3).*

This has the following interpretation: given zero and first-order data, the only constraint on the second-order data of an initial configuration of surfaces satisfying ‘order one compatibility at $t = 0$ ’ is condition (1) on the mean curvatures at the junction curve, while (3) specifies how the mean curvatures determine the normal velocity of the interface Γ_t , in particular at $t = 0$ (since the normal velocity $\dot{a} \cdot n$ is independent of how we parametrize Γ_t , we denote it by $\dot{\Gamma}_n$):

$$\dot{\Gamma}_n = \frac{v_E}{v^3 \sqrt{3}}(H^1 - H^2) - \frac{\partial_n w^3}{v^3} H^3.$$

Geometrically in $\mathbb{R}^3$, condition (3) says that the component V^{nor} (normal to the junction Λ_t) of the velocity vector, which is independent of how one parametrizes Λ_t , is determined by the H^I via:

$$V^{nor} = H^3 N^3 + \mu^3 T^3 = H^3 N^3 + \frac{1}{\sqrt{3}}(H^1 - H^2)T^3.$$

Proof. This is computational and elementary, so we just describe the main steps. One just has to observe that (as for curves), the constraints on first derivatives arising from writing $N^1 + N^2 = N^3$ componentwise can be ‘solved’, in the sense of the following elementary algebraic lemma:

Lemma 1.2. Given real numbers $\alpha > 0$ and a^1, a^2, a^3 , let $v^I = \sqrt{\alpha^2 + (a^I)^2}$; suppose these numbers satisfy the following relations:

$$\frac{1}{v^1} + \frac{1}{v^2} = \frac{1}{v^3}, \quad \frac{a^1}{v^1} + \frac{a^2}{v^2} = \frac{a^3}{v^3}.$$

Then (assuming $a^2 > a^1$ to normalize the labeling):

$$\begin{aligned} \frac{1}{v^1} &= \frac{1}{v^3} \left(\frac{1}{2} + \frac{\sqrt{3}}{2} \frac{a^3}{\alpha} \right), & \frac{1}{v^2} &= \frac{1}{v^3} \left(\frac{1}{2} - \frac{\sqrt{3}}{2} \frac{a^3}{\alpha} \right), \\ \frac{a^1}{v^1} &= \frac{1}{2} \frac{a^3}{v^3} - \frac{\sqrt{3}}{2} \frac{\alpha}{v^3}, & \frac{a^2}{v^2} &= \frac{1}{2} \frac{a^3}{v^3} + \frac{\sqrt{3}}{2} \frac{\alpha}{v^3}. \end{aligned}$$

Remark. This lemma has the following equivalent geometric formulation: let $\omega^1, \omega^2, \omega^3$ be vectors in $\mathbb{R}^2$ of the same length, satisfying $\omega^1 + \omega^2 = \omega^3$. Then, denoting by R_θ the counterclockwise rotation operator in $\mathbb{R}^2$ (by θ radians), we have:

$$\omega^1 = R_{\pi/3}[\omega^3], \quad \omega^2 = R_{5\pi/3}[\omega^3].$$

Since n is the inner unit normal of $D_t^1 = D_t^2$, the condition $a^2 > a^1$ corresponds to labeling the surfaces so that Σ^2 is below Σ^1 near Λ_t .

To prove proposition 1.1, we substitute the values of v^I and $d_n w^I / v^I$ given by lemma 1.2 (for $I = 1, 2$, in terms of the same quantities for $I = 3$) into relations (2)-(6). Using relation (1), we find that each of (2) and (4) is equivalent to (3). Where $d_\tau w \neq 0$, adding and subtracting the relations obtained from (5) and (6) with this substitution, we recover respectively (1) and (3), completing the proof.

There is no ‘order 1’ compatibility condition to consider at the fixed boundary $\partial\Omega$, since the Neumann condition $d_\omega w^3 = 0$ is first-order.

2. An equivalent system on a fixed domain.

In this section we describe new dependent variables $u^I(x, t), \tilde{\rho}(x, t)$ defined on fixed cylindrical domains $D_0^I \times [0, T]$, $\Omega \times [0, T]$ (resp.), and a parabolic system with conjugation conditions, equivalent to the original system for $w^I(y, t)$ on $D_t^I, t \in [0, T]$.

2.1. An extension operator. $\Gamma_0 \subset \Omega \subset \mathbb{R}^2$ is a simple closed curve of class $C^{3+\alpha}$, bounding domains $D_0^1 = D_0^2$ on the inside and $D_0^3 = \Omega - \bar{D}_0^1$;

denote by ν the inner unit normal vector field of Γ_0 (of class $C^{2+\alpha}$) and by $r : \mathcal{N}_0 \rightarrow \mathbb{R}$ the oriented distance to Γ_0 , where $\mathcal{N}_0$ is a fixed tubular neighborhood of Γ_0 . We have $r \in C^{3+\alpha}(\mathcal{N}_0)$, and the natural extension $\nu = Dr$ of ν to $\mathcal{N}_0$. With $\bar{r} := \sup_{\mathcal{N}_0} r$, consider the smaller tubular neighborhood:

$$\mathcal{N} := \{x \in \mathcal{N}_0; |r(x)| < r_0 := \min\{\bar{r}/2, 1/2\}\}.$$

Denoting by $\hat{\zeta} \in C^\infty(\mathbb{R}; [0, 1])$ a fixed function supported in $[-r_0, r_0]$, equal to 1 in $[-r_0/2, r_0/2]$ with $|\hat{\zeta}'| \leq 3/r_0$ everywhere, we see that $\zeta(x)$, defined as $\hat{\zeta} \circ r$ in $\mathcal{N}$ and identically zero outside of $\mathcal{N}$, is in $C^{3+\alpha}(\mathbb{R}^2)$ and coincides with r in $\mathcal{N}_1 := r^{-1}(-r_0/2, r_0/2) \subset \mathcal{N}$.

Using ζ we extend $\nu = Dr$ from $\mathcal{N}_1$ to $\mathbb{R}^2$, by setting $\bar{\nu} := \zeta \nu \in C^{2+\alpha}(\mathbb{R}^2; \mathbb{R}^2)$. $\bar{\nu}$ is supported in $\mathcal{N}$ and satisfies $|\bar{\nu}| \leq 1$ in $\mathbb{R}^2$. Note that $D_\nu \nu \equiv 0$ in $\mathcal{N}_1$, while:

$$D_{\bar{\nu}} \bar{\nu} = \zeta D_\nu (\zeta \nu) = \zeta (d_\nu \zeta) \nu = (d_\nu \zeta) \bar{\nu},$$

which is supported in $\mathcal{N} - \mathcal{N}_1$.

We will need a fixed extension operator $C^{k+\alpha}(\Gamma_0) \rightarrow C^{k+\alpha}(\mathbb{R}^2)$, defined in a standard way using ζ (for $k \geq 2$). Given $\rho \in C^{k+\alpha}(\Gamma_0)$, extend ρ to $\mathcal{N}_0$ by projection along normal line segments (so $d_\nu \rho \equiv 0$ in $\mathcal{N}_0$), then set:

$$\tilde{\rho} := \begin{cases} \zeta \rho, & x \in \mathcal{N}_0 \\ 0, & x \in \mathbb{R}^2 - \mathcal{N} \end{cases}$$

It is clear that $\tilde{\rho}$ is supported in $\mathcal{N}$, and that $\|\tilde{\rho}\|_{C^{k+\alpha}(\mathbb{R}^2)} \leq c_\zeta \|\rho\|_{C^{k+\alpha}(\Gamma_0)}$, where c_ζ depends only on ζ and on $\|r\|_{C^{3+\alpha}(\mathcal{N}_0)}$. Since the extension operator is t -independent, the same estimate holds for parabolic Hölder norms when ρ depends on t . Note also:

$$\begin{aligned} D_{\bar{\nu}}(\tilde{\rho} \bar{\nu}) &= \zeta \nu(\tilde{\rho}) \bar{\nu} + \tilde{\rho} D_{\bar{\nu}} \bar{\nu} \\ &= \zeta(\nu(\zeta) \rho + \zeta \nu(\rho)) \bar{\nu} + \tilde{\rho} (d_\nu \zeta) \bar{\nu} \\ &= \zeta((d_\nu \zeta) \rho + 0) \bar{\nu} + \tilde{\rho} (d_\nu \zeta) \bar{\nu} \\ &= 2(d_\nu \zeta) \tilde{\rho} \bar{\nu}, \end{aligned}$$

supported in $\mathcal{N} - \mathcal{N}_1$.

If U is already defined on a set containing Γ_0 , we denote by $\mathcal{E}[U]$ the restriction-extension operator:

$$\mathcal{E}[U] := (U|_{\Gamma_0})^\sim.$$

2.2. Diffeomorphism. For $t > 0$ small enough, the interface Γ_t will be contained in $\mathcal{N}_1$, and we may parametrize it as a normal graph over Γ_0 :

$$\Gamma_t = \{x + \rho(x, t)\nu(x); x \in \Gamma_0\}, \text{ for some } \rho : \Gamma_0 \times [0, T] \rightarrow \mathbb{R}.$$

Recall the subscript t always denotes ‘function (or map, or set) at time t ’, so $\rho_t(x) := \rho(x, t)$. We use the extension operator $\mathcal{E}$ to define a diffeomorphism φ_t of $\mathbb{R}^2$ (or of Ω), for $t \in [0, T]$:

$$\varphi_t(x) = x + \tilde{\rho}_t(x)\bar{\nu}(x), \quad \varphi_0 = id.$$

For $t \in [0, T]$ small enough, φ_t is a diffeomorphism (of class $C^{2+\alpha}$), equal to the identity in $\mathbb{R}^2 - \mathcal{N}$, and mapping:

$$\Omega \rightarrow \Omega, \quad \Gamma_0 \rightarrow \Gamma_t, \quad D_0^I \rightarrow D_t^I, \quad I = 1, 2, 3,$$

diffeomorphically in each case. If $\rho \in C^{2+\alpha, 1+\alpha/2}(\Gamma_0 \times [0, T])$, φ is in the same Hölder class as a function of (x, t) . The differential of φ_t is given by:

$$D\varphi_t = \mathbb{I} + (D\tilde{\rho}_t)\bar{\nu} + \tilde{\rho}_t D\bar{\nu};$$

in particular, in light of the calculation in 2.1, $D\varphi_t$ maps $\bar{\nu}$ via:

$$D\varphi_t[\bar{\nu}] = \bar{\nu} + D\bar{\nu}(\tilde{\rho}_t\bar{\nu}) = [1 + 2(d_\nu\zeta)\tilde{\rho}_t]\bar{\nu},$$

which implies:

$$(D\varphi_t)^{-1}[\bar{\nu}] = \frac{1}{1 + 2(d_\nu\zeta)\tilde{\rho}_t}\bar{\nu} := \bar{\zeta}(\tilde{\rho})\bar{\nu}.$$

Here $\bar{\zeta}(\tilde{\rho}) \in C^{3+\alpha}(\mathbb{R}^2)$, and is identically 1 in $\mathcal{N}_1 \cap (\mathbb{R}^2 - \mathcal{N})$, with modulus bounded above by 1 everywhere (provided only we have $|\tilde{\rho}_t| \leq r_0/12$ for $t \in [0, T]$). In particular, on Γ_0 :

$$D\varphi_t|_{\Gamma_0} : \nu(x) \mapsto \nu(x + \rho_t(x)\nu(x))$$

(recall $\nu := Dr$ on $\mathcal{N}_0$). Of course, $\nu(x + \rho_t(x)\nu(x))$ is in general not normal to Γ_t .

2.3 Equations over D_0^I . Assume $w_t^I : D_t^I \rightarrow \mathbb{R}$ are solutions of $(MCM)_w$ for $t \in [0, T]$. Define, for $I = 1, 2, 3$:

$$u^I(x, t) = w^I(\varphi_t(x), t), x \in D_0^I; \quad w^I(y, t) = u^I(\psi_t(y), t), y \in D_t^I$$

(where $\psi_t := (\varphi_t)^{-1}$). A standard calculation yields (omitting the superscript I sometimes, and using the Einstein summation convention throughout):

$$0 = \partial_t w - g^{ij}(Dw)D_{i,j}^2 w = \partial_t u - h^{ab}D_{a,b}^2 u + (\partial_c u)(\partial_t \psi^c - g^{ij}D_{i,j}^2 \psi^c),$$

or equivalently:

$$\partial_t u - h^{ab}D_{a,b}^2 u - (\partial_c u)(\partial_i \psi^c)(\partial_t \varphi^i - h^{ab}\partial_{a,b}^2 \varphi^i) = 0,$$

where $h_{ab} = \partial_a \varphi^i \partial_b \varphi^j g_{ij}$ is the pullback metric (of g under φ). Since the parametrization over D_0^I of Σ^I is:

$$F_t^I(x) = G_t^I(\varphi_t(x)) = [\varphi_t(x), w_t^I(\varphi_t(x))] = [\varphi_t(x), u_t^I(x)] \in \mathbb{R}^2 \times \mathbb{R},$$

we have for $x \in D_0^I$:

$$\begin{aligned} h_{ab}^I &= \langle \partial_a F_t^I, \partial_b F_t^I \rangle = \partial_a \varphi_t \cdot \partial_b \varphi_t + (\partial_a u_t^I)(\partial_b u_t^I) \\ &= \delta_{ab} + \partial_a(\tilde{\rho}_t \bar{\nu}) \cdot \partial_b(\tilde{\rho}_t \bar{\nu}) + (\partial_a u_t^I)(\partial_b u_t^I) \\ &:= \delta_{ab} + h'_{ab}(\tilde{\rho}_t, D\tilde{\rho}_t, Du_t^I). \end{aligned}$$

Define the operator on functions (or maps) $f(x, t)$ on $D_0^I \times [0, T]$:

$$L_{h^I}[f] := \partial_t f - tr_{h^I} D^2 f.$$

The equation on D_0^I is:

$$L_{h^I}[u^I] - Du^I \cdot (D\varphi_t)^{-1}[L_{h^I}\varphi] = 0.$$

We may express $L_h[\varphi]$ in terms of $\tilde{\rho}$ and $\bar{\nu}$:

$$L_h[\varphi] = L_h[\tilde{\rho}]\bar{\nu} - \tilde{\rho} tr_h D^2 \bar{\nu} - 2\langle D\tilde{\rho}, D\bar{\nu} \rangle_h.$$

Using $(D\varphi)^{-1}[\bar{\nu}] = \bar{\zeta}(\tilde{\rho})\bar{\nu}$, we find:

$$(D\varphi)^{-1}[L_h\varphi] = \bar{\zeta}(\tilde{\rho})L_h[\tilde{\rho}]\bar{\nu} - (D\varphi)^{-1}[\tilde{\rho} tr_h D^2 \bar{\nu} + 2\langle D\tilde{\rho}, D\bar{\nu} \rangle_h].$$

Hence the system in the variables $(u^I, \tilde{\rho})$ is:

$$L_{h^I}[u^I] - \bar{\zeta}(\tilde{\rho})(d_{\bar{\nu}} u^I) L_{h^I}[\tilde{\rho}] + Du^I \cdot (D\varphi_t)^{-1}[\tilde{\rho} tr_{h^I} D^2 \bar{\nu} + 2\langle D\tilde{\rho}, D\bar{\nu} \rangle_{h^I}] = 0, \quad (PDE)_{(u^I, \rho)}$$

where:

$$h_{ab}^I = \delta_{ab} + h'_{ab}(\tilde{\rho}, D\tilde{\rho}, Du^I), \quad D\varphi_t = \mathbb{I} + D(\tilde{\rho}\bar{\nu}),$$

$$\langle D\tilde{\rho}, D\bar{\nu} \rangle_{h^I} = (h^I)^{ab}(\partial_a \tilde{\rho})(\partial_b \bar{\nu}) \in \mathbb{R}^2.$$

2.4. Boundary conditions at Γ_0 . (More precisely, these are ‘conjugation conditions’, since Γ_0 is an ‘internal boundary’ in Ω). We begin by computing how $D\varphi_t|_{\Gamma_0}$ acts on the orthonormal frame $\{\nu, \tau_0\}$, $\tau_0 := -\nu^\perp$ (where $\perp$ denotes counterclockwise $\pi/2$ rotation in $\mathbb{R}^2$). Parametrizing Γ_0 by arc length s , we have:

$$\Gamma'_0(s) = \tau_0, \quad \nu' = -k_0\tau_0$$

(with k_0 the curvature of Γ_0). Composing with φ_t , we obtain the parametrization of Γ_t : $\Gamma_t(s) = \Gamma_0(s) + \rho_t(\Gamma_0(s))\nu(s)$, with tangent and inner unit normal vectors:

$$\Gamma'_t(s) = (1 - k_0\rho)\tau_0 + (d_{\tau_0}\rho)\nu, \quad n = \frac{(1 - k_0\rho)\nu - (d_{\tau_0}\rho)\tau_0}{[(1 - k_0\rho)^2 + (d_{\tau_0}\rho)^2]^{1/2}}.$$

Using $D_\nu\nu = 0, d_\nu\rho = 0, D_{\tau_0}\nu = \nu'$ on Γ_0 , we find for an arbitrary $v \in \mathbb{R}^2$ and $x \in \Gamma_0$:

$$D\varphi_t(x)[v] = v + (v \cdot \tau_0)(d_{\tau_0}\rho)\nu - (v \cdot \tau_0)(k_0\rho)\tau_0,$$

in particular verifying again:

$$D\varphi_t[\nu] = \nu, \quad D\varphi_t[\tau_0] = (1 - k_0\rho)\tau_0 + (d_{\tau_0}\rho)\nu = \Gamma'_t(s).$$

We need vectors mapping to n and to $\tau := -n^\perp$ under $D\varphi_t$. It is easy to see that, defining:

$$\begin{aligned} \hat{\tau}_0 &:= [(1 - k_0\rho)^2 + (d_{\tau_0}\rho)^2]^{-1/2}\tau_0 := a_{11}\tau_0, \\ \mu &= (1 - k_0\rho)^{-1}\{[(1 - k_0\rho)^2 + (d_{\tau_0}\rho)^2]^{1/2}\nu - (d_{\tau_0}\rho)\hat{\tau}_0\} \\ &:= a_{21}\tau_0 + a_{22}\nu, \end{aligned}$$

we have for these vector fields $\mu = \mu(\rho, d_{\tau_0}\rho), \hat{\tau}_0 = \hat{\tau}_0(\rho, d_{\tau_0}\rho)$:

$$D\varphi_t(x)[\mu] = n, \quad D\varphi_t(x)[\hat{\tau}_0] = \tau, \quad x \in \Gamma_0.$$

(The expressions with the a_{ij} are used in section 4.) Hence, for $x \in \Gamma_0$:

$$d_n w^I(\varphi_t(x)) = d_\mu u^I(x); \quad d_\tau w^I(\varphi_t(x)) = d_{\hat{\tau}_0} u^I(x),$$

$$v^I = [1 + (d_\tau w^I)^2 + (d_n w^I)^2]^{1/2} = [1 + (d_{\hat{\tau}_0} u^I)^2 + (d_\mu u^I)^2]^{1/2} \\ := \mathcal{G}(\rho, d_{\tau_0} \rho, d_{\tau_0} u^I, d_\nu u^I).$$

(*Remark:* It is easy to see that $\mu - \nu = -(d_{\tau_0} \rho) \tau_0 + O(\rho^2 + |d_{\tau_0} \rho|^2)$; at $t = 0$, $\mu \equiv \nu$ on γ_0 , since $\rho \equiv 0$.)

We now state the conjugation conditions in terms of u^I, ρ :

$$\begin{cases} u^1(x, t) = u^2(x, t) = u^3(x, t) & (BC^0)_{(u^I, \rho)} \\ \frac{1}{v^1} + \frac{1}{v^2} = \frac{1}{v^3} & (BC^1)_{(u^I, \rho)} \\ \frac{d_\nu u^1}{v^1} + \frac{d_\nu u^2}{v^2} = \frac{d_\nu u^3}{v^3} & (BC^2)_{(u^I, \rho)} \end{cases}$$

In the computation of the second angle condition, we use the fact that $d_{\hat{\tau}_0} u^I$ is independent of I on Γ_0 (which follows from the matching condition). Here $v^I = \mathcal{G}(\rho, d_{\tau_0} \rho, d_{\tau_0} u^I, d_\nu u^I)$ has the expression given above.

Since the diffeomorphism φ_t is the identity on $\partial\Omega$, we have the additional (Neumann) boundary condition for u^3 :

$$\partial_\omega u^3 = 0 \text{ on } \partial\Omega.$$

(Recall $\partial D_0^3 = \Gamma_0 \sqcup \partial\Omega$.)

3. The transformed system.

The system described in the preceding section includes 4 scalar conjugation conditions, but only three evolution equations for the four unknowns u^I, ρ ; there is no explicit evolution equation for ρ (or its extension $\tilde{\rho}$). Yet (assuming the evolution of triple junctions is well-posed as a PDE) it is clear geometrically that the evolution of the u^I determines that of ρ . In this section, following the technique introduced in the paper [BaconneauLunardi] we show it is possible to define new dependent variables U^I on $D_0^I \times [0, T]$ in such a way that $\tilde{\rho}$ is recovered from the U^I via the extension operator $\mathcal{E}$. The drawback (as in [BaconneauLunardi]) is having to introduce ‘non-local highest-order terms’ in the resulting equation for the U^I .

Following [BaconneauLunardi], define $U^I : D_0^I \times [0, T] \rightarrow \mathbb{R}$ via:

$$u^I(x, t) = u_0^I(x) + (d_\nu u_0^I)(x) \tilde{\rho}(x, t) + U^I(x, t).$$

(Note $U_{|t=0}^I \equiv 0$.) The matching conditions $(BC^0)_{(u^I, \rho)}$ on Γ_0 imply:

$$\rho d_\nu u_0^1 + U^1 = \rho d_\nu u_0^2 + U^2 = \rho d_\nu u_0^3 + U^3 \text{ on } \Gamma_0.$$

Equivalently, we have the following equalities on Γ_0 :

$$\rho|_{\Gamma_0} = \frac{U^2 - U^1}{d_\nu u_0^1 - d_\nu u_0^2} = \frac{U^3 - U^2}{d_\nu u_0^2 - d_\nu u_0^3} = \frac{U^3 - U^1}{d_\nu u_0^1 - d_\nu u_0^3}.$$

The last equality may be written in the form:

$$\frac{U^3 - U^2}{U^3 - U^1} = \frac{d_\nu u_0^2 - d_\nu u_0^3}{d_\nu u_0^1 - d_\nu u_0^3} = -\frac{v_0^2}{v_0^1},$$

which we regard as a new form of the matching condition. We used the fact that the angle conditions at $t = 0$ imply:

$$\frac{1}{v_0^1}(d_\nu u_0^1 - d_\nu u_0^3) = \frac{1}{v_0^2}(d_\nu u_0^3 - d_\nu u_0^2).$$

This new matching condition may also be written in the form:

$$\frac{U^3 - U^2}{v_0^2} + \frac{U^3 - U^1}{v_0^1} = 0,$$

or equivalently (using the first angle condition at $t = 0$):

$$\frac{U^1}{v_0^1} + \frac{U^2}{v_0^2} = \frac{U^3}{v_0^3} \text{ on } \Gamma_0 \quad (BC^0)_{U^I},$$

which we adopt as the matching condition for the U^I system.

The main point of the method is that the relation:

$$\rho = \delta_0(U^2 - U^1) \text{ on } \Gamma_0, \quad \delta_0 := (d_\nu u_0^1 - d_\nu u_0^2)^{-1}$$

allows us to express $\tilde{\rho}$ in terms of $\mathcal{E}[U^2 - U^1]$.

To explain how this is done, we take a moment to examine the extension operator applied to the product of two functions, f, g defined on Γ_0 . Denote by the subscript *rad* the extension to $\mathcal{N}_0$ constant along normals of a function defined on Γ_0 . By definition, on $\mathcal{N}_0$ we have:

$$(fg)^\sim = \zeta(fg)_{rad} = (f)_{rad}\tilde{g}$$

(this is true on $\mathcal{N}$, and both sides vanish on $\mathcal{N}_0 - \mathcal{N}$). Let χ be a smooth function in $\mathbb{R}^2$, equal to 1 on $\mathcal{N}$ and vanishing outside $\mathcal{N}_0$. Given f on Γ_0 , define $\hat{f}$ on $\mathbb{R}^2$ by:

$$\hat{f} = \chi f_{rad} \text{ on } \mathcal{N}_0, \quad \hat{f} = 0 \text{ on } \mathbb{R}^2 - \mathcal{N}_0.$$

Then it is easy to check that:

$$(fg)^{\sim} = \hat{f}\tilde{g} \text{ on } \mathbb{R}^2.$$

Thus we may recover $\tilde{\rho}$ from $U^2 - U^1$ as follows:

$$\tilde{\rho} = \hat{\delta}_0 \mathcal{E}[U^2 - U^1].$$

We now express the angle boundary conditions in terms of the U^I . Since:

$$u^I = u_0^I + \delta_0(U^2 - U^1)d_\nu u_0^I + U^I \text{ on } \Gamma_0,$$

we have on Γ_0 :

$$\begin{aligned} d_{\tau_0} u^I &= d_{\tau_0} u_0^I + d_{\tau_0}[\delta_0(U^2 - U^1)d_\nu u_0^I] + d_{\tau_0} U^I \\ &:= A_{tan}(U^2 - U^1, d_{\tau_0}(U^2 - U^1)) + d_{\tau_0} U^I \end{aligned}$$

(A_{tan} for ‘affine’, as a function of the indicated arguments.) For the normal derivatives, we use the expression defining U^I on D_0^I , and recall that $d_\nu \tilde{\rho} = 0$ on Γ_0 ; hence (using also $D_\nu \nu = 0$):

$$\begin{aligned} d_\nu u_{|\Gamma_0}^I &= d_\nu u_0^I + \delta_0(U^2 - U^1)D^2 u_0^I(\nu, \nu) + d_\nu U^I \\ &:= A_{nor}(U^2 - U^1) + d_\nu U^I. \end{aligned}$$

In section 2 we found functional expressions $v_{|\Gamma_0}^I = \mathcal{G}(\rho, d_{\tau_0} \rho, d_{\tau_0} u^I, d_\nu u^I)$. If in the expressions defining the vector fields $\hat{\tau}_0$ and μ on Γ_0 we replace ρ and $d_{\tau_0} \rho$ by $\delta_0(U^2 - U^1)$ and $d_{\tau_0}[\delta_0(U^2 - U^1)]$ (resp.) and use the expressions just found for $d_{\tau_0} u^I$ and $d_\nu u^I$, we obtain:

$$v_{|\Gamma_0}^I = \mathcal{G}(\delta_0(U^2 - U^1), d_{\tau_0}[\delta_0(U^2 - U^1)], A_{tan}(U^2 - U^1, d_{\tau_0}(U^2 - U^1)) + d_{\tau_0} U^I, A_{nor}(U^2 - U^1) + d_\nu U^I).$$

From now on we adopt the notational convention for the summation of quantities depending on a superscript $I = 1, 2, 3$:

$$\sum_I' a^I := a^1 + a^2 - a^3.$$

It follows from the above (using also the boundary conditions for the u_0^I) that the boundary conditions take the form:

$$\begin{cases} \sum_I' \frac{U^I}{v_0^I} = 0 & (BC^0)_{U^I} \\ \sum_I' \frac{1}{v^I} = 0 & (BC^1)_{U^I} \\ \sum_I' \frac{d_\nu U^I}{v^I} + \sum_I' \frac{d_\nu u_0^I}{v^I} + \delta_0(U^2 - U^1) \sum_I' \frac{D^2 u_0^I(\nu, \nu)}{v^I} = 0 & (BC^2)_{U^I} \end{cases}$$

Computing the equation satisfied by the U^I is straightforward. First we write down the expression for $L_{h^I}[u^I]$, isolating the lower-order terms:

$$L_{h^I}[u^I] = L_{h^I}[U^I] + (d_{\bar{\nu}} u_0^I) L_{h^I}[\tilde{\rho}] + \mathcal{L}_{h^I}^{(1)I}(\tilde{\rho}, D\tilde{\rho}),$$

where:

$$\mathcal{L}_{h^I}^{(1)I}(\tilde{\rho}, D\tilde{\rho}) := -2\langle D\tilde{\rho}, D(d_{\bar{\nu}} u_0^I) \rangle_{h^I} - \text{tr}_{h^I} D^2 u_0^I - \tilde{\rho} \text{tr}_{h^I} D^2(d_{\bar{\nu}} u_0^I).$$

Now using the equation $(PDE)_{(u^I, \rho)}$ from section 2 and the definitions:

$$\mathcal{C}_{h^I}(\tilde{\rho}, D\tilde{\rho}) := (D\varphi)^{-1}[\tilde{\rho} \text{tr}_{h^I} D^2 \bar{\nu} + 2\langle D\tilde{\rho}, D\bar{\nu} \rangle_{h^I}] \text{ with } D\varphi = \mathbb{I} + D(\tilde{\rho} \bar{\nu}),$$

$$\mathcal{L}_{h^I}^{(2)I}(\tilde{\rho}, D\tilde{\rho}) := D(u_0^I + \tilde{\rho} d_{\bar{\nu}} u_0^I) \cdot \mathcal{C}_{h^I},$$

we find:

$$\begin{aligned} & L_{h^I}[U^I] + [(d_{\bar{\nu}} u_0^I - \bar{\zeta}(\tilde{\rho}) d_{\bar{\nu}}(u_0^I + \tilde{\rho} d_{\bar{\nu}} u_0^I + U^I))] L_{h^I}[\tilde{\rho}] \\ & + DU^I \cdot \mathcal{C}_{h^I}(\tilde{\rho}, D\tilde{\rho}) + \mathcal{L}_{h^I}^{(1)I}(\tilde{\rho}, D\tilde{\rho}) + \mathcal{L}_{h^I}^{(2)I}(\tilde{\rho}, D\tilde{\rho}) = 0. \end{aligned}$$

(Note that $(D\varphi)^{-1}$ depends on $D\tilde{\rho}$, and that the first-order terms involving the metric h^I depend on DU^I , in addition to $\tilde{\rho}$ and $D\tilde{\rho}$.)

This may be regarded as a quasilinear system in U^I , if we agree to replace $\tilde{\rho}$ by $\hat{\delta}_0 \mathcal{E}[U^2 - U^1]$ at every occurrence of $\tilde{\rho}$ or $D\tilde{\rho}$. Due to the presence of the restriction-extension operator $\mathcal{E}$, this introduces ‘non-local terms’, even to highest order.

To make the dependence on U^I a bit more explicit, compute:

$$\begin{aligned} & L_{h^I}[\tilde{\rho}] = L_{h^I}[\hat{\delta}_0 \mathcal{E}[U^2 - U^1]] \\ & = \hat{\delta}_0 L_{h^I}[\mathcal{E}[U^2 - U^1]] - 2\langle D\hat{\delta}_0, D(\mathcal{E}[U^2 - U^1]) \rangle_{h^I} - \mathcal{E}[U^2 - U^1] \text{tr}_{h^I} D^2 \hat{\delta}_0. \end{aligned}$$

Thus we have the equation for U^I :

$$L_{h^I}[U^I] + \mathcal{A}(\tilde{\rho}, D\tilde{\rho}, DU^I)|_{\tilde{\rho}=\hat{\delta}_0 \mathcal{E}[U^2-U^1]} L_{h^I}[\mathcal{E}[U^2 - U^1]]$$

$$= \mathcal{F}_U^I(\tilde{\rho}, D\tilde{\rho}, \mathcal{E}[U^2 - U^1], D(\mathcal{E}[U^2 - U^1]))|_{\tilde{\rho}=\hat{\delta}_0\mathcal{E}[U^2-U^1]},$$

where:

$$\begin{aligned} \mathcal{A}(\tilde{\rho}, D\tilde{\rho}, DU^I) &= \hat{\delta}_0[(d_{\tilde{\nu}}u_0^I - \bar{\zeta}(\tilde{\rho})d_{\tilde{\nu}}(u_0^I + \tilde{\rho}d_{\tilde{\nu}}u_0^I + U^I)]; \\ \mathcal{F}_U^I &= 2\langle D\hat{\delta}_0, D(\mathcal{E}[U^2 - U^1]) \rangle_{h^I} + \mathcal{E}[U^2 - U^1]tr_{h^I}D^2\hat{\delta}_0 \\ &\quad - D(u_0^I + \tilde{\rho}d_{\tilde{\nu}}u_0^I + U^I) \cdot \mathcal{C}_{h^I} \\ &\quad + 2\langle D\tilde{\rho}, D(d_{\tilde{\nu}}u_0^I) \rangle_{h^I} + tr_{h^I}D^2u_0^I + \tilde{\rho}tr_{h^I}D^2(d_{\tilde{\nu}}u_0^I), \end{aligned}$$

and $\mathcal{C}_{h^I}$ was given above.

We record this in abbreviated form (changing the notation slightly to exhibit the dependence of h^I and $\mathcal{A}$ on U):

$$L_{h_U^I}[U^I] + \mathcal{A}_U^I L_{h_U^I}[\mathcal{E}[U^2 - U^1]] = \mathcal{F}_U^I \quad (PDE)_{U^I}$$

The critical term is $\mathcal{A}_U^I L_{h_U^I}[\mathcal{E}[U^2 - U^1]]$, due to the ‘non-local dependence’ on U^I (i.e., it is not given by the action of a differential operator on the U^I , so we may not quote standard linear parabolic theory at this point). This term has the important property:

$$\mathcal{A}_{|t=0}^I \equiv 0 \text{ on } D_0^I.$$

This is easy to see from the definition, since $\mathcal{A}^I$ is $\hat{\delta}_0$ multiplied by the expression:

$$d_{\tilde{\nu}}u_0^I - \bar{\zeta}(\tilde{\rho})d_{\tilde{\nu}}(u_0^I + \tilde{\rho}d_{\tilde{\nu}}u_0^I + U^I).$$

At $t = 0$ we have $U^I \equiv 0$, $\tilde{\rho} \equiv 0$ (since $\rho \equiv 0$) and therefore $\bar{\zeta}(\tilde{\rho}) \equiv 1$. This makes it possible to treat this linear system as a small perturbation of a standard linear parabolic system, as long as T is small. This was also the general strategy adopted in [BaconneauLunardi], but we actually deal with this term in a different way (in a sense, less sophisticated; see Section 4). $\tilde{\rho}_{|t=0} = 0$ also implies $\mathcal{C}_{h^I} = 0$ at $t = 0$.

Finally, we record the dependence of the metric h^I on DU^I :

$$\begin{aligned} h_{ab}^I &= \delta_{ab} + \partial_a(\tilde{\rho}\tilde{\nu}) \cdot \partial_b(\tilde{\rho}\tilde{\nu}) \\ &\quad + (\partial_a u_0^I + \partial_a(\tilde{\rho}d_{\tilde{\nu}}u_0^I) + \partial_a U^I)(\partial_b u_0^I + \partial_b(\tilde{\rho}d_{\tilde{\nu}}u_0^I) + \partial_b U^I), \end{aligned}$$

where we set $\tilde{\rho} = \hat{\delta}_0\mathcal{E}[U^2 - U^1]$.

Note also the two properties of the term $\mathcal{F}_U^I$: it involves derivatives of U^I up to first order, and equals $tr_{h_0^I} D^2 u_0^I = v_0^I H_0^I$ at $t = 0$.

4. The associated linear system. In this section we describe the (standard) fixed-point argument used to prove local existence, based on the solution of an associated linear system with ‘non-local terms’.

4.1 Linearization of the conjugation conditions.

The angle conjugation conditions $(BC)_{U^I}$ involve sums of $1/v^I$, so we begin by considering the linearization $\mathcal{L}$ of $1/v$ at points of Γ_0 , where:

$$v|_{\Gamma_0} = \sqrt{1 + (d_\mu u)^2 + (d_{\hat{\tau}_0} u)^2} = v(\rho, d_{\tau_0} \rho, DU)$$

and we linearize at $\rho \equiv 0, U \equiv 0$. We find:

$$\mathcal{L}\left(\frac{1}{v}\right) = -\frac{1}{v_0^3} [(d_\nu u_0) \mathcal{L}(d_\mu u) + (d_{\tau_0} u_0) \mathcal{L}(d_{\hat{\tau}_0} u)]$$

From the expressions for $\hat{\tau}_0$ and μ given in section 2:

$$\hat{\tau}_0 = a_{11} \tau_0, \quad \mu = a_{21} \tau_0 + a_{22} \nu, \quad a_{ij}|_{\rho \equiv 0} = \delta_{ij},$$

we have:

$$\mathcal{L}(d_{\hat{\tau}_0} u) = d_{\tau_0} U + (d_\nu u_0) d_{\tau_0} \rho + \mathcal{L}(a_{11}) d_{\tau_0} u_0 + d^2 u_0(\tau_0, \nu) \rho,$$

$$\mathcal{L}(d_\mu u) = d_\nu U + \mathcal{L}(a_{21}) d_{\tau_0} u_0 + \mathcal{L}(a_{22}) d_\nu u_0 + d^2 u_0(\nu, \nu) \rho.$$

Straightforward computation from the expressions for the a_{ij} given in section 2 shows that:

$$\mathcal{L}(a_{11}) = k_0 \rho, \quad \mathcal{L}(a_{21}) = -d_{\tau_0} \rho, \quad \mathcal{L}(a_{22}) = 0.$$

Substituting in the expression for $\mathcal{L}(v^{-1})$, we find that the coefficient of $d_{\tau_0} \rho$ vanishes, yielding:

$$\mathcal{L}\left(\frac{1}{v}\right) = -\frac{1}{v_0^3} \{Du_0 \cdot DU + [d_\nu u_0 d^2 u_0(\nu, \nu) + d_{\tau_0} u_0 d^2 u_0(\nu, \tau_0) + (d_{\tau_0} u_0)^2 k_0] \rho\}$$

Restoring the superscript I , we write this (always at Γ_0) in the form:

$$\frac{1}{v^I} = \frac{1}{v_0^I} - \frac{1}{(v_0^I)^3} Du_0^I \cdot DU^I + \gamma_0^I \rho + E^{(1)I}(\rho, d_{\tau_0} \rho, DU^I)$$

where:

$$\gamma_0^I = -\frac{1}{(v_0^I)^3} [d_\nu u_0^I d^2 u_0^I(\nu, \nu) + d_{\tau_0} u_0^I d^2 u_0^I(\nu, \tau_0) + (d_{\tau_0} u_0^I)^2 k_0]$$

and $|E^{(1)I}| = O(\rho^2 + (d_{\tau_0} \rho)^2 + |DU^I|^2)$, with constants depending only on the initial data.

The first angle condition $(BC^1)_{U^I}$ then becomes, with $E^{(1)} = \sum_I' E^{(1)I}$:

$$\sum_I' \frac{Du_0^I}{(v_0^I)^3} \cdot DU^I - \rho \sum_I' \gamma_0^I = E^{(1)}(\rho, d_{\tau_0} \rho, DU).$$

Now make the substitution $\rho \rightarrow \delta_0(U^2 - U^1)$, valid on Γ_0 . We find:

$$\sum_I' \frac{Du_0^I}{(v_0^I)^3} \cdot DU^I + \gamma_0^{(1)}(U^2 - U^1) = E^{(1)}(\rho, d_{\tau_0} \rho, DU)|_{\rho=\delta_0(U^2-U^1)},$$

where:

$$\gamma_0^{(1)} := -\delta_0 \sum_I' \gamma_0^I.$$

The second angle condition $(BC^2)_{U^I}$ has the form:

$$\sum_I' \frac{d_\nu U^I}{v^I} + \sum_I' \frac{d_\nu u_0^I}{v^I} + \rho \sum_I' \frac{D^2 u_0^I(\nu, \nu)}{v_0^I} = 0,$$

or (using the linearization of v^{-1} computed above):

$$\sum_I' \left(\frac{1}{v_0^I} d_\nu U^I - \frac{d_\nu u_0^I}{(v_0^I)^3} Du_0^I \cdot DU^I \right) + \left(\sum_I' (d_\nu u_0^I) \gamma_0^I + \frac{d^2 u_0^I(\nu, \nu)}{v_0^I} \right) \rho = E^{(2)},$$

where $|E^{(2)}(\rho, d_{\tau_0} \rho, DU)| = O(\rho^2 + (d_{\tau_0} \rho)^2 + |DU|^2)$. Again making the substitution $\rho \rightarrow \delta_0(U^2 - U^1)$, we find:

$$\sum_I' \left(\frac{1}{v_0^I} d_\nu U^I - \frac{d_\nu u_0^I}{(v_0^I)^3} Du_0^I \cdot DU^I \right) + \gamma_0^{(2)}(U^2 - U^1) = E^{(2)}(\rho, d_{\tau_0} \rho, DU)|_{\rho=\delta_0(U^2-U^1)},$$

where:

$$\gamma_0^{(2)} = \delta_0 \sum_I' \left((d_\nu u_0^I) \gamma_0^I + \frac{d^2 u_0^I(\nu, \nu)}{v_0^I} \right).$$

In summary, the conjugation conditions may be written in linearized form:

$$\begin{cases} \sum_I ' \frac{U^I}{v_0^I} = 0 & (LBC^0)_{U^I} \\ \sum_I ' B_0^{(1)I} \cdot DU^I + \gamma_0^{(1)}(U^2 - U^1) = E^{(1)}(U, DU) & (LBC^1)_{U^I} \\ \sum_I ' B_0^{(2)I} \cdot DU^I + \gamma_0^{(2)}(U^2 - U^1) = E^{(2)}(U, DU) & (LBC^2)_{U^I} \end{cases}$$

(with zero initial conditions.) Here we defined:

$$B_0^{(1)I} := \frac{Du_0^I}{(v_0^I)^3}, \quad B_0^{(2)I} := \frac{1}{v_0^I} \nu - \frac{d_\nu u_0^I}{(v_0^I)^3} Du_0^I.$$

Observe (for future use) that, since $1 - (d_\nu u_0)^2/(v_0)^2 = [1 + (d_{\tau_0} u_0)^2]/(v_0)^2$, we have:

$$\sum_I ' B_0^{(2)I} \cdot DU^I = [1 + (d_{\tau_0} u_0)^2] \sum_I ' \frac{1}{(v_0^I)^3} d_\nu U^I - (d_{\tau_0} u_0) \sum_I ' \frac{d_\nu u_0^I}{(v_0^I)^3} d_{\tau_0} U^I.$$

(The notation uses the fact that $d_{\tau_0} u_0^I$ is independent of I at points of Γ_0).

4.2 The fixed-point scheme.

We write the quasilinear system $(PDE)_{U^I}/(BC)_{U^I}$ in a slightly modified ‘linearized form’, as follows:

$$(LPDE/LBC)_U \begin{cases} L_{h_0^I}[U^I] = f_U^I(x, t) & \text{on } D_0^I \times (0, T] \\ \sum_I ' \frac{U^I}{v_0^I} = 0, \quad \mathbb{B}_0[DU] + \gamma_0[U] = E_U & \text{on } \Gamma_0, \quad d_\omega U|_{\partial\Omega}^3 = 0 \end{cases}$$

where:

$$f_U^I(x, t) := \mathcal{F}_U^I + [(h_0^I)^{ab} - (h_U^I)^{ab}] \partial_{a,b}^2 U^I - \mathcal{A}_U^I L_{h_U^I}[\mathcal{E}[U^2 - U^1]]$$

and we collapsed two linearized angle conditions into one, introducing a slight change in notation. $h_0^I = g_0^I$ is the induced metric at $t = 0$.

This will be solved by a standard fixed-point argument in the space:

$$\mathbb{X}_{R,T} := \{U = (U^1, U^2, U^3); U^I \in C^{2+\alpha, 1+\alpha/2}(D_0^I \times [0, T]), U|_{t=0} \equiv 0\},$$

with suitable choices of R and T , to be described soon. (We organize the fixed-point argument in the same way as [BaconneauLunardi].)

Define an operator $\Phi : V \mapsto U$ by assigning to $V \in \mathbb{X}_{R,T}$ the solution of the linear system:

$$(LPDE_0) \left\{ \begin{array}{l} L_{h_0^I}[U^I] = f_V^I(x, t) \quad \text{on } D_0^I \times (0, T] \\ \sum_I' \frac{U^I}{v_0^I} = 0, \quad \mathbb{B}_0[DU] + \gamma_0[U] = E_V \text{ on } \Gamma_0, \quad d_\omega U|_{\partial\Omega}^3 = 0 \end{array} \right.$$

Consider the following assumptions for system $(LPDE_0)$:

(1) (Regularity of coefficients) $h_0^I \in C^\alpha(D_0^I)$, $\mathbb{B}_0, \gamma_0 \in C^{1+\alpha}(\Gamma_0)$, $(v_0^I)^{-1} \in C^{2+\alpha}(\Gamma_0)$;

(2) (Complementarity) The conjugation operator $(\sum_I' \frac{U^I}{v_0^I}, \mathbb{B}_0[DU])$ satisfies the ‘complementarity conditions’ on Γ_0 , with respect to the operators $L_{h_0^I}$.

Under these assumptions, it is a classical fact that this system (with vanishing initial data) has a unique solution, provided $f_V^I \in C^{\alpha, \alpha/2}(D_0^I \times [0, T])$ and $E_V \in C^{1+\alpha, (1+\alpha)/2}(\Gamma_0 \times [0, T])$ satisfy also:

(3) (Compatibility with $U|_{t=0}^I \equiv 0$.)

$$\sum_I' \frac{f_V^I}{v_0^I}|_{t=0} = 0, \quad E_V|_{t=0} = 0.$$

The solution satisfies the estimate:

$$\|U^I\|_{2+\alpha} \leq M_0 \left(\sum_I \|f_V^I\|_\alpha + \|E_V\|_{1+\alpha} \right),$$

with M_0 depending only on the Hölder norms of the coefficients in (1).

The verification of condition (1) under the assumption $u_0^I \in C^{3+\alpha}(D_0^I)$ is straightforward. Verifying (3) is also easy, since when $U^I \equiv 0$ we have $E_U = 0$ and $f_U^I = \text{tr}_{h_0^I} D^2 u_0^I$, so that the compatibility condition amounts to:

$$\sum_I' \frac{1}{v_0^I} \text{tr}_{h_0^I} D^2 u_0^I = 0,$$

which is just the compatibility condition considered in section 1 (equivalent to $\sum_I' H_0^I = 0$ on Γ_0 .)

Complementarity (assumption (2)) also holds for the system $(LPDE_0)$, but verifying this is more technical- it is done in the next section. Assuming (2) for the moment, we conclude the map Φ is well-defined.

4.3 Contraction estimates in Hölder norms.

To finish the argument, we must verify that (with suitable choices of R and T) Φ maps into $\mathbb{X}_{T,R}$ and is a contraction in this space.

Let $V_1, V_2 \in \mathbb{X}_{R,T}$. Then $W = \Phi(V_1) - \Phi(V_2)$ is the solution of the linear problem:

$$(LPDE_W) \begin{cases} L_{h_0^I}[W^I] = f_{V_1}^I - f_{V_2}^I & \text{on } D_0^I \times (0, T] \\ \sum_I' \frac{W^I}{v_0^I} = 0, \quad \mathbb{B}_0[DW] + \gamma_0[W] = E_{V_1} - E_{V_2} & \text{on } \Gamma_0, \quad d_\omega W|_{\partial\Omega}^3 = 0 \end{cases}$$

(with zero initial conditions.) Thus we have the estimate:

$$\|W^I\|_{2+\alpha} \leq M_0 \left(\sum_I \|f_{V_1}^I - f_{V_2}^I\|_\alpha + \|E_{V_1} - E_{V_2}\|_{1+\alpha} \right).$$

We write the difference $f_{V_1}^I - f_{V_2}^I$ as the sum of five terms:

$$\begin{aligned} f_{V_1}^I - f_{V_2}^I &= \mathcal{F}_{V_1}^I - \mathcal{F}_{V_2}^I \\ &+ (h_{V_2}^{Iab} - h_{V_1}^{Iab}) \partial_{ab}^2 V_1^I + (h_0^{Iab} - h_{V_2}^{Iab}) \partial_{ab}^2 (V_1^I - V_2^I) \\ &+ (\mathcal{A}_{V_1}^I - \mathcal{A}_{V_2}^I) L_{h_{V_1}^I} [\mathcal{E}(V_1^I)] \\ &+ \mathcal{A}_{V_2}^I (h_{V_1}^{Iab} - h_{V_2}^{Iab}) \partial_{ab}^2 V_1^I \\ &+ \mathcal{A}_{V_2}^I h_{V_2}^{Iab} \partial_{ab}^2 (V_1^I - V_2^I) \\ &:= f^{(1)I} + \dots + f^{(5)I}. \end{aligned}$$

We need bounds for each of the $f^{(i)I}$ in terms of $\|V_1 - V_2\|_{2+\alpha}$, R and T . The details are lengthy but standard, and it suffices to state the estimates with a brief justification. (c_0 denotes a constant depending only on the initial data, which may change from one occurrence to the next.)

$$\|\mathcal{F}_{V_1}^I - \mathcal{F}_{V_2}^I\|_\alpha \leq c_0 T^{\alpha/2} \|V_1 - V_2\|_{2+\alpha},$$

since $\mathcal{F}_V^I$ depends only on $V, DV, \mathcal{E}(V)$ and $D(\mathcal{E}(V))$, and vanishes at $t = 0$. (The ‘nonlocality’ of $\mathcal{E}$ does not make the estimates harder, since this operator is linear in V and bounded in $C^{2+\alpha, 1+\alpha/2}$ norm.)

$$\|f^{(2)I}\|_\alpha \leq c_0 T^{\alpha/2} \|V_1 - V_2\|_{2+\alpha} R + c_0 T^{\alpha/2} \|V_1 - V_2\|_{2+\alpha},$$

since $h_{V_2}^I - h_{V_1}^I$ and $h_0^I - h_{V_2}^I$ depend only on derivatives of V_1, V_2 up to first order and vanish at $t = 0$.

$$(\mathcal{A}_{V_1}^I - \mathcal{A}_{V_2}^I) L_{h_{V_1}^I} [\mathcal{E}(V_1^I)] \leq c_0 T^{\alpha/2} \|V_1 - V_2\|_{2+\alpha} R,$$

since $\mathcal{A}_{V_1}^I - \mathcal{A}_{V_2}^I$ depends on derivatives of V_1, V_2 only up to first order and vanishes at $t = 0$.

$$\mathcal{A}_{V_2}^I (h_{V_1}^{Iab} - h_{V_2}^{Iab}) \partial_{ab}^2 V_1^I \leq c_0 T^{\alpha/2} \|V_1 - V_2\|_{2+\alpha} R^2,$$

since $\mathcal{A}_{V_2}^I$ depends on derivatives of V_2 up to first order, and vanishes at $t = 0$.

Finally,

$$\mathcal{A}_{V_2}^I h_{V_2}^{Iab} \partial_{ab}^2 (V_1^I - V_2^I) \leq c_0 T^{\alpha/2} \|V_1 - V_2\|_{2+\alpha} R^2,$$

for the same reason.

A bit more involved (but also needed) is the estimate over Γ_0 :

$$\|E_{V_1} - E_{V_2}\|_{1+\alpha} \leq c_0 T^{\alpha/2} \|V_1 - V_2\|_{2+\alpha} R.$$

Although the $1 + \alpha$ norm of E_V involves $D^2 V$, the estimate holds since E_V is of quadratic order in V and DV .

We conclude that Φ is a $1/2$ contraction, provided we choose R and T so that $16c_0(1 + R^2)T^{\alpha/2} \leq \frac{1}{2}$ ($16 = 5 \times 3 + 1$). Assuming this inequality holds, applying the contraction estimate to the case $V_2 = 0$ yields, for any $V \in \mathbb{X}_{R,T}$:

$$\|\Phi(V)\|_{2+\alpha} \leq \|\Phi(0)\|_{2+\alpha} + \frac{1}{2}R,$$

where $\Phi(0)$ is the solution (with zero initial data) of the system $(LPDE)_0$ with data $f_V^I = tr_{h_0^I} D^2 u_0$, $E_V = 0$, satisfying the estimate:

$$\|\Phi(0)\|_{2+\alpha} \leq M_0 \|tr_{h_0^I} D^2 u_0^I\|_{\alpha}.$$

The choices of R and T are made as follows. As observed earlier, if the C^2 geometry of the configuration at time t is sufficiently close to that at $t = 0$, all the geometric constructions used make sense; in particular the operator $\mathcal{E}$ and $\mathcal{F}_U^I$ are well defined. Choose $r > 0$ so that $\|U^I\|_{C^2(D_0^I)} < r$ quantifies ' u^I sufficiently close to u_0^I '. Then pick $T_0 > 0$ so that $U|_{t=0} \equiv 0$

and $\|U\|_{2+\alpha} < T_0$ imply $\|U^I\|_{C^2(D_0^I)} < r$. Now choose $R > 0$ sufficiently large to ensure:

$$M_0 \|tr_{h_0^I} D^2 u_0^I\|_\alpha < \frac{1}{2}R.$$

Finally, given R we pick $T < T_0$ small enough that $16c_0(1 + R^2)T^{\alpha/2} \leq \frac{1}{2}$ holds.

With these choices we guarantee, on the one hand, that Φ is a contraction; and, on the other, that:

$$\|\Phi(V)\|_{2+\alpha} \leq \|\Phi(0)\|_{2+\alpha} + \frac{1}{2}R < R$$

for any $V \in \mathbb{X}_{R,T}$. Thus Φ maps into $\mathbb{X}_{R,T}$, and has a unique fixed point U . This fixed point is the unique solution of $(LPDE)_U/(LBC)_U$ with zero initial data. This concludes the proof of local existence for the system $(PDE)_{(u^I, \bar{\rho})}$, and hence for the original system $(PDE)_{w^I}$. (Except for the verification of complementarity for $(LPDE)_0$, carried out in the next section).

5. The complementarity condition.

In this section we verify that the system of conjugation conditions along Γ_0 satisfies the ‘complementarity conditions’ (Lopatinski-Shapiro) with respect to the 3×3 linear parabolic operator $L_{g_0^I}[U^I]$. (We use the conditions as stated in [EidelmanZhitarasu].) Explicitly, the operator is:

$$\begin{aligned} L_{g_0^I} U^I &= \partial_t U^I - g_0^{Iab} \partial_{ab}^2 U^I \text{ on } \Omega \times [0, T], \\ g_{0ab}^I &= \delta_{ab} + \partial_a u_0^I \partial_b u_0^I, \quad g_0^{Iab} = \delta_{ab} - \frac{\partial_a u_0^I \partial_b u_0^I}{(v_0^I)^2}, \\ v_0^I &= \sqrt{1 + (d_{\tau_0} u_0^I)^2 + (d_\nu u_0^I)^2} \text{ on } \Gamma_0. \end{aligned}$$

The conjugation operator has three components:

$$\begin{cases} B_0[Y] = \sum_I ' \frac{U^I}{v_0^I} \\ B_1[Y] = \sum_I ' \frac{n_0^I}{(v_0^I)^2} d_\nu U^I + \lambda_0 \sum_I ' \frac{1}{(v_0^I)^2} d_{\tau_0} U^I, \\ B_2[Y] = (1 + \lambda_0^2) \sum_I ' \frac{1}{(v_0^I)^3} d_\nu U^I - \lambda_0 \sum_I ' \frac{n_0^I}{(v_0^I)^3} d_{\tau_0} U^I, \end{cases}$$

where in this section we adopt the notation:

$$n_0^I := d_\nu u_0^I|_{\Gamma_0}, \quad \lambda_0 := d_{\tau_0} u_0^I|_{\Gamma_0}.$$

Fix $x_0 \in \Gamma_0$ and ‘straighten the boundary’ via a diffeomorphism χ from a neighborhood of $0 \in \mathbb{R}^2$ (with coordinates $z = (z_1, z_2)$) to a neighborhood of x_0 in Ω , and with the mapping properties:

$$\chi : \{z_1 = 0\} \rightarrow \Gamma_0, \quad \chi(0) = x_0, \quad d\chi(0) : \partial_{z_1} \mapsto \nu(x_0), \quad d\chi(0) : \partial_{z_2} \mapsto \tau_0(x_0).$$

Let $Y^I(z, t) = U^I(\chi(z), t)$. From this point on the symbols γ^I , n_0^I , λ_0 , v_0^I will denote the values of the corresponding functions at the fixed point $x_0 \in \Gamma_0$.

The transformed operator is:

$$L_0[Y^I] := \partial_t Y^I - \gamma^{Iij} \partial_{z_i z_j}^2 Y^I.$$

$\gamma^I = \chi(0)^* g_0^I(x_0)$ is the pullback metric tensor, with components:

$$\gamma_{11}^I = 1 + (n_0^I)^2, \quad \gamma_{12}^I = \lambda_0 n_0^I, \quad \gamma_{22}^I = 1 + \lambda_0^2,$$

and inverse:

$$\gamma^{I11} = \frac{1 + \lambda_0^2}{(v_0^I)^2}, \quad \gamma^{I12} = -\frac{\lambda_0 n_0^I}{(v_0^I)^2}, \quad \gamma^{I22} = \frac{1 + (n_0^I)^2}{(v_0^I)^2}.$$

In the new coordinates, the components of the conjugation operator are:

$$\begin{cases} B_0[Y] = \sum_I ' \frac{Y^I}{v_0^I} \\ B_1[Y] = \sum_I ' \frac{n_0^I}{(v_0^I)^2} \partial_{z_1} Y^I + \lambda_0 \sum_I ' \frac{1}{(v_0^I)^2} \partial_{z_2} Y^I, \\ B_2[Y] = (1 + \lambda_0^2) \sum_I ' \frac{1}{(v_0^I)^3} \partial_{z_1} Y^I - \lambda_0 \sum_I ' \frac{n_0^I}{(v_0^I)^3} \partial_{z_2} Y^I, \end{cases}$$

The next step is to consider the Fourier-Laplace transform $\hat{Y}^I(\tau, \xi, p)$ of $Y^I(z_1, z_2, t)$: Fourier transform in z_2 corresponds to the variable $\xi \in \mathbb{R}$, while Laplace transform in t corresponds to $p \in \mathbb{C}$. Furthermore, we adjust the signs so that $\{\tau > 0\}$ corresponds to $D_0^1 = D_0^2$ (that is, $\tau = z_1$ for $\hat{Y}^1$ and $\hat{Y}^2$) and also to D_0^3 (that is, $\tau = -z_1$ for $\hat{Y}^3$). (This is how we deal with the fact that we have a conjugation problem, rather than a boundary-value problem.) This introduces a sign σ^I in the transformed operator ($\sigma^1 = \sigma^2 = 1, \sigma^3 = -1$), which is an ordinary differential operator in the variable τ , for each fixed (p, ξ) :

$$\gamma^{I11} \frac{d^2 \hat{Y}^I}{d\tau^2} + 2\sigma^I i\xi \gamma^{I12} \frac{d\hat{Y}^I}{d\tau} - \xi^2 \gamma^{I22} \hat{Y}^I - p \hat{Y}^I.$$

We are interested in solutions of the corresponding ODE which decay as $\tau \rightarrow +\infty$. If we assume the exponential form:

$$\hat{Y}_{\xi,p}^I(\tau) = e^{i\tau\rho_I} \hat{Y}_{\xi,p}^I(0),$$

this corresponds to $\mathbb{I}m(\rho_I) > 0$. The $\rho_I \in \mathbb{C}$ depend on ξ and p , and are roots of the indicial equation:

$$\gamma^{I11}\rho_I^2 + 2\sigma^I\gamma^{I12}\xi\rho_I + \gamma^{I22}\xi^2 + p = 0.$$

Using the expressions given earlier for the γ^{Iij} , we find for the roots:

$$\begin{aligned} \rho_I &= \frac{1}{\gamma^{I11}}[-\sigma^I\gamma^{I12}\xi + i\sqrt{\xi^2 \det[\gamma^I]^{-1} + p\gamma^{I11}}] \\ &= \frac{v_0^I}{1 + \lambda_0^2}[\sigma^I\lambda_0\frac{n_0^I}{v_0^I}\xi + i\sqrt{\Delta}], \end{aligned}$$

where $\Delta = \xi^2 + p(1 + \lambda_0^2)$ and we take the branch $\mathbb{R}e(\sqrt{\Delta}) > 0$. The boundary operators for the ODE are obtained from the correspondence:

$$\partial_{z_1} Y_{|z_1=0}^I \longrightarrow \sigma^I \frac{d\hat{Y}^I}{d\tau} \Big|_{\tau=0} = \sigma^I i\rho_I \hat{Y}^I(0), \quad \partial_{z_2} Y_{|z_1=0}^I \longrightarrow i\xi \hat{Y}^I(0).$$

We find:

$$\begin{aligned} B_0[Y] &\longrightarrow \sum_I \frac{\hat{Y}^I(0)}{v_0^I} \\ B_1[Y] &\longrightarrow \sum_I \left[\frac{n_0^I}{(v_0^I)^3} \sigma^I (i\rho_I) + \frac{\lambda_0}{(v_0^I)^3} (i\xi) \right] \hat{Y}^I(0) \\ B_2[Y] &\longrightarrow \sum_I \left[\frac{1+\lambda_0^2}{(v_0^I)^3} \sigma^I (i\rho_I) - \frac{\lambda_0 n_0^I}{(v_0^I)^3} (i\xi) \right] \hat{Y}^I(0) \end{aligned}$$

The complementarity condition is the statement that, for each (p, ξ) in the set:

$$\mathcal{A} := \{(p, \xi) \in \mathbb{C} \times \mathbb{R}; |p| + |\xi| > 0, (1 + \lambda_0^2)\mathbb{R}e(p) > -\xi^2\}$$

the map $\hat{Y}^I(0) \mapsto B_I[Y]$ defines an isomorphism of $\mathbb{C}^3$. In matrix form (after canceling a common factor $(v_0^I)^{-1}$ in each column) we are interested in:

$$\mathbb{T}_{(p,\xi)} := \begin{bmatrix} 1 & 1 & -1 \\ \frac{n_0^1}{(v_0^1)^2}\rho_1 + \frac{\lambda_0}{(v_0^1)^2}\xi & \frac{n_0^2}{(v_0^2)^2}\rho_2 + \frac{\lambda_0}{(v_0^2)^2}\xi & \frac{n_0^3}{(v_0^3)^2}\rho_3 - \frac{\lambda_0}{(v_0^3)^2}\xi \\ \frac{1+\lambda_0^2}{(v_0^1)^2}\rho_1 - \frac{\lambda_0 n_0^1}{(v_0^1)^2}\xi & \frac{1+\lambda_0^2}{(v_0^2)^2}\rho_2 - \frac{\lambda_0 n_0^2}{(v_0^2)^2}\xi & \frac{1+\lambda_0^2}{(v_0^3)^2}\rho_3 + \frac{\lambda_0 n_0^3}{(v_0^3)^2}\xi \end{bmatrix}$$

Lemma 5.1. For each $(p, \xi) \in \mathcal{A}$, $\det(\mathbb{T}_{(p, \xi)}) \neq 0$.

Proof. We begin by introducing some notation. Consider the vectors $t_I \in \mathbb{R}^2, r_I \in \mathbb{C}^2$:

$$t_I = \frac{1}{v_0^I} [n_0^I, \sqrt{1 + \lambda_0^2}], \quad r_I = \frac{1}{v_0^I} [\sqrt{1 + \lambda_0^2} \rho_I, \lambda_0 \sigma^I \xi].$$

Then:

$$\frac{n_0^1 \rho_1}{v_0^1 v_0^1} + \frac{\lambda_0}{v_0^1} \frac{\xi}{v_0^1} = \frac{1}{\sqrt{1 + \lambda_0^2}} t_1 \cdot r_1, \quad \frac{1 + \lambda_0^2 \rho_1}{v_0^1 v_0^1} - \frac{n_0^1 \lambda_0 \xi}{v_0^1 v_0^1} = t_1 \cdot r_1^\perp,$$

and analogously for the other two columns. (Here we use the ‘real dot product’, i.e. $t_I \cdot r_I = t_I \cdot \operatorname{Re}(r_I) + it_I \cdot \operatorname{Im}(r_I)$).

With this notation, we find (after cancelation of common factors and using $t_I \cdot r_I^\perp = -t_I^\perp \cdot r_I$) we must show that the matrix:

$$\begin{bmatrix} 1 & 1 & -1 \\ t_1 \cdot r_1 & t_2 \cdot r_2 & t_3 \cdot r_3 \\ t_1^\perp \cdot r_1 & t_2^\perp \cdot r_2 & t_3^\perp \cdot r_3 \end{bmatrix}$$

has non-zero determinant **det**. Expanding, we have:

$$\mathbf{det} = (t_1^\perp \cdot r_1)(t_2 \cdot r_2) - (t_2^\perp \cdot r_2)(t_1 \cdot r_1) - (t_1 \cdot r_1)(t_3^\perp \cdot r_3) + (t_1^\perp \cdot r_1)(t_3 \cdot r_3) + (t_2 \cdot r_2)(t_3^\perp \cdot r_3) - (t_2^\perp \cdot r_2)(t_3 \cdot r_3).$$

The argument has two parts. In the first part, we use the symmetries in the conjugation conditions, encoded in the properties of t_I :

$$t_1 + t_2 = t_3, \quad |t_I| = 1.$$

From the geometric version of lemma 1.2, we can express t_1 and t_2 in terms of t_3 :

$$t_1 = R_{\pi/3} t_3 = \frac{1}{2} t_3 + \frac{\sqrt{3}}{2} t_3^\perp, \quad t_2 = R_{5\pi/3} t_3 = \frac{1}{2} t_3 - \frac{\sqrt{3}}{2} t_3^\perp.$$

Substituting these values in the expression for **det**, we obtain:

$$\begin{aligned} \mathbf{det} &= \frac{\sqrt{3}}{2} (r_1 + r_2) \cdot [(t_3 \cdot r_3) t_3 + (t_3^\perp \cdot r_3) t_3^\perp] + \frac{1}{2} (r_1 - r_2) \cdot [(t_3^\perp \cdot r_3) t_3 - (t_3 \cdot r_3) t_3^\perp] \\ &\quad + r_2 \cdot \left\{ \frac{\sqrt{3}}{2} [(r_1 \cdot t_3) t_3 + (r_1 \cdot t_3^\perp) t_3^\perp] + \frac{1}{2} [(t_3^\perp \cdot r_1) t_3 - (t_3 \cdot r_1) t_3^\perp] \right\}, \end{aligned}$$

which simplifies to give an expression in terms of the r_I only:

$$\mathbf{det} = \frac{\sqrt{3}}{2}(r_1 \cdot r_3 + r_2 \cdot r_3 + r_1 \cdot r_2) + \frac{1}{2}(r_1 \cdot r_2^\perp + r_2 \cdot r_3^\perp + r_3 \cdot r_1^\perp).$$

In the second part of the argument, we use symmetries in the r_I . Recalling the expressions for the roots ρ_I , we have (with $a_I := \operatorname{Re}(r_I)$, $b_I := \operatorname{Im}(r_I)$, both in $\mathbb{R}^2$):

$$r_I = a_I + ib_I, \quad a_I = \lambda_0 \xi \left[\frac{1}{\sqrt{1 + \lambda_0^2}} \frac{\sigma^I n_0^I}{v_0^I}, \frac{\sigma^I}{v_0^I} \right], \quad b_I = \frac{\sqrt{\Delta}}{\sqrt{1 + \lambda_0^2}} [1, 0].$$

We see that $b_I := b$ is independent of I , while the a_I satisfy:

$$\sum_I a_I = 0, \quad |a_I|^2 = \frac{\lambda_0^2 \xi^2}{1 + \lambda_0^2} := |a|^2$$

(independent of I). Consider first the imaginary part of $\mathbf{det}$. We have:

$$\operatorname{Im}(r_I \cdot r_J) = \operatorname{Re}(r_I) \operatorname{Im}(r_J) + \operatorname{Re}(r_J) \operatorname{Im}(r_I) = (a_I + a_J) \cdot b,$$

and from this it follows easily that $\operatorname{Im}(r_1 \cdot r_3 + r_2 \cdot r_3 + r_1 \cdot r_2) = 0$. Similarly, since $r_I^\perp = a_I^\perp + ib^\perp$, we have:

$$\operatorname{Im}(r_I \cdot r_J^\perp) = a_I \cdot b^\perp + a_J^\perp \cdot b,$$

and again we have: $\operatorname{Im}(r_1 \cdot r_2^\perp + r_2 \cdot r_3^\perp + r_3 \cdot r_1^\perp) = 0$. Thus $\operatorname{Im}(\mathbf{det}) = 0$.

Turning to the real part, since $\operatorname{Im}(r_I) \cdot \operatorname{Im}(r_I)^\perp = b \cdot b^\perp = 0$:

$$\operatorname{Re}(r_I \cdot r_J) = a_I \cdot a_J - |b|^2, \quad \operatorname{Re}(r_I \cdot r_J^\perp) = a_I \cdot a_J^\perp - b \cdot b^\perp = a_I \cdot a_J^\perp.$$

Using the version of lemma 1.2 for the case $\sum_I a_I = 0, |a_I| = |a|$:

$$a_1 = R_{4\pi/3} a_3, \quad a_2 = R_{2\pi/3} a_3$$

we see that:

$$\begin{aligned} a_1 \cdot a_3 &= a_2 \cdot a_3 = a_1 \cdot a_2 = -\frac{1}{2}|a|^2, \\ a_1 \cdot a_2^\perp &= a_2 \cdot a_3^\perp = a_3 \cdot a_1^\perp = \frac{\sqrt{3}}{2}|a|^2. \end{aligned}$$

We conclude:

$$\mathbf{det} = \frac{\sqrt{3}}{2}(-\frac{3}{2}|a|^2 - 3|b|^2) + \frac{1}{2}\frac{3\sqrt{3}}{2}|a|^2 = -(3\sqrt{3}/2)|b|^2 = -(3\sqrt{3}/2)\frac{|\Delta|}{1 + \lambda_0^2}.$$

This is non-zero for any $(p, \xi) \in \mathcal{A}$ (since $\Re(\Delta) > 0$ in $\mathcal{A}$), as we had to show.

Remark. Note that this calculation depends on the fact that the boundary conditions (that is, the t_I) and the roots of the characteristic equation (that is, the r_I , and ultimately the coefficients of the operator, which come from the induced metric at the junction) satisfy the same type of symmetry. That is, complementarity of this particular set of conjugation conditions seems to be linked to the fact that we are dealing with mean curvature motion; this suggests that local well-posedness may fail for a more general parabolic system (with the same conjugation conditions.)

References

- [BaconneauLunardi] Baconneau, O.; Lunardi, A. *Smooth solutions to a class of free boundary parabolic problems*. Trans. Amer. Math. Soc. **356** (2004), no. 3, 987–1005.
- [BronsardReitich] Bronsard, Lia; Reitich, Fernando On three-phase boundary motion and the singular limit of a vector-valued Ginzburg-Landau equation. Arch. Rational Mech. Anal.**124** (1993), no. 4, 355–379.
- [Dierkes et al.] U.Dierkes, S.Hildebrandt A.Küster, O.Wohlrab, *Minimal Surfaces 1*. Grundlehren der Mathematischen Wissenschaften v.295, Springer-Verlag 1992. ISBN 3-540-53169-6
- [EidelmanZhitarasu] Eidelman, S. D.; Zhitarashu, N. V. *Parabolic boundary value problems*. Operator Theory: Advances and Applications, 101. Birkhuser Verlag, Basel, 1998. xii+298 pp. ISBN: 3-7643-2972-6
- [Freire] Freire,A. *Mean Curvature Motion of Graphs with Constant Contact Angle and Moving Boundaries* (preprint, May 2008-arXiv:0805.4592)
- [LensSeminar] *Evolution of convex lens-shaped networks under curve shortening flow* O. Schnürer, A. Azouani, M. Georgi, J. Hell, N. Jangle, A. Koeller, T. Marxen, S. Ritthaler, M. Sáez, F. Schulze, B. Smith, (Lens Seminar, FU Berlin 2007), arXiv:0711.1108
- [Mantegazza et al.] Mantegazza,C., Novaga,M., Tortorelli,V., *Motion by curvature of planar networks*. Ann. Sc. Norm. Super. Pisa Cl. Sci. **5**(3) (2004) no. 2, 235–324.